arXiv:1210.2743v1 [math.NT] 9 Oct 2012

CHARACTER SUMS DETERMINED BY LOW DEGREE ISOGENIES OF ELLIPTIC CURVES

DUSTIN MOODY AND CHRISTOPHER RASMUSSEN

Abstract. We consider character sums determined by isogenies of elliptic curves over finite fields. We prove a congruence condition for character sums attached to arbitrary cyclic isogenies, and produce explicit formulas for isogenies of degree $m \leq 8$.

1. Introduction

Let $p > 3$ be prime, and let h_p denote the class number of $\mathbb{Q}(\sqrt{-p})$. Let h_p^* be either 0 or h_p, as p is congruent to 1 or 3 modulo 4, respectively. A well-known consequence of Dirichlet's analytic class number formula is the following identity:

$$-\frac{1}{p}\sum_{x=1}^{p-1} x\left(\frac{x}{p}\right) = h_p^*. \tag{1}$$

In [4], a similar class number identity is proved, where the sum runs over the $\mathbb{F}_p$ points of an elliptic curve, and the quadratic residue character is replaced by the cokernel character of a 2-isogeny of elliptic curves. More precisely:

Let k be a field with $\mathrm{char}(k) \neq 2$, and let E_1/k be the elliptic curve:

$$E_1 : y^2 = x^3 + ax^2 + bx, \qquad b \neq 0, a^2 \neq 4b, \tag{2}$$

which has a k-rational 2-isogeny given by

$$\tau(x, y) = \left(\frac{x^2 + ax + b}{x}, y\frac{(b - x^2)}{x^2}\right), \tag{3}$$

to the curve

$$E_2 : x^3 - 2ax^2 + (a^2 - 4b)x.$$

For $k = \mathbb{F}_p$, define a character χ_τ on E_2 by

$$\chi_\tau(P) = \begin{cases} 1 & P \in \tau\big(E_1(\mathbb{F}_p)\big) \\ -1 & P \notin \tau\big(E_1(\mathbb{F}_p)\big) \end{cases}.$$

Then for any prime $p > 3$ of good reduction, the quantity

$$S_{\tau,p} = \sum_{\substack{P \in E_2(\mathbb{F}_p) \\ P \neq \infty}} \{x_P - a\} \chi_\tau(P) \tag{4}$$

satisfies the congruence $S_{\tau,p} \equiv 0 \pmod{p}$. Furthermore, $-\frac{1}{p} S_{\tau,p} - h_p^*$ is bounded in absolute value by a constant C_τ independent of p. The notation $\{x\}$ denotes the lifting from $\mathbb{F}_p$ to $\mathbb{Z}$ which satisfies $0 \leq \{x\} \leq (p-1)$ for all $x \in \mathbb{F}_p$.

In this paper, we extend the study to isogenies of degree greater than 2. In Section 2, we deduce an exact sequence related to the isogeny formulas of Vélu over finite fields. In Section 3 we generalize the character sum from [4] to higher degree isogenies, and prove a relation that transports character sums across a normalized isogeny. This generalizes the congruence property of $S_{\tau,p}$. In Section 4, we apply the results to determine explicit formulas for families of isogenies of degree $m \leq 8$.

2. Isogenies

2.1. **Vélu's formula.** If the kernel of a separable isogeny ϕ is finite with cardinality d, then ϕ is referred to as a d-isogeny, and the isogeny is said to have degree d. It is known that if d is composite, then every d-isogeny can be decomposed into a composition of prime degree isogenies. For this reason, we mainly concern ourselves with prime degree isogenies. We let $\hat{\phi}$ denote the dual isogeny to ϕ, which satisfies $\phi \circ \hat{\phi} = [d]$, $\hat{\phi} \circ \phi = [d]$. Here $[d]$ is the multiplication by d map.

Let E be an elliptic curve defined over a field k. Given a finite k-rational subgroup $\mathcal{F} \leq E$, let E' denote the quotient $E/\mathcal{F}$. Vélu has given formulas [8] for both a Weierstrass equation for E', and a separable isogeny $\phi\colon E \to E'$ whose kernel is exactly $\mathcal{F}$. For a point $P = (x_P, y_P) \in E - \mathcal{F}$, the formula is:

$$\phi(P) = \left(x_P + \sum_{Q \in \mathcal{F} - \{\infty\}} (x_{P+Q} - x_Q),\ y_P + \sum_{Q \in \mathcal{F} - \{\infty\}} (y_{P+Q} - y_Q) \right). \tag{5}$$

2.2. **Normalization.** In this subsection, we fix a field k, not of characteristic 2 or 3. Let E/k be an elliptic curve. We may write a Weierstrass equation for E in the form $y^2 = x^3 + ax + b$. Let E'/k be another elliptic curve, and let $\phi\colon E \to E'$ be a separable isogeny defined over k. If the pullback of the invariant differential ω' of E' along ϕ is equal to the invariant differential ω of E, then ϕ is said to be *normalized.* As the space of differentials is one dimensional, we know $\phi^*\omega' = c_\phi \omega$, for some $c_\phi \in k^*$. If $c_\phi = 1$, then $\phi^*\omega' = \omega$ and ϕ is normalized.

We collect a few results about normalized isogenies which we will need for the proof of Theorem 6. Gonzalez [1] noted some of these, but only included a proof of Proposition 4. We include proofs for completeness. We believe that these results are well-known, but not all collected in one place in the literature.

Lemma 1. *Let E_1/k, E_2/k be elliptic curves, and let $\phi_1 \colon E_1 \to E_2$ and $\phi_2 \colon E_1 \to E_2$ be k-rational isogenies. Then $c_{\phi_1+\phi_2} = c_{\phi_1} + c_{\phi_2}$.*

Proof. Let ω_i denote the invariant differential of E_i. This is an immediate consequence of the fact that $(\phi_1 + \phi_2)^*\omega_2 = \phi_1^*\omega_2 + \phi_2^*\omega_2$ ([6, Thm. III.5.2]). Since $\phi_i^*\omega_2 = c_{\phi_i}\omega_1$, we have $c_{\phi_1+\phi_2} = c_{\phi_1} + c_{\phi_2}$. ☐

Lemma 2. *Let E_i/k $(i = 1, 2, 3)$ be elliptic curves, and let $\phi_1 \colon E_1 \to E_2$, $\phi_2 \colon E_2 \to E_3$ be k-rational isogenies. Then $c_{\phi_2 \circ \phi_1} = c_{\phi_1} c_{\phi_2}$.*

Proof. Let ω_i denote the invariant differential of E_i. Then

$$(\phi_2 \circ \phi_1)^*\omega_3 = \phi_1^*\phi_2^*\omega_3 = \phi_1^*(c_{\phi_2}\omega_2) = c_{\phi_1} c_{\phi_2} \omega_1,$$

and the result follows. ☐

Corollary 3. *The composition of two normalized isogenies is normalized.*

Proposition 4. *Let E_1/k and E_2/k be elliptic curves, and let ϕ_1, $\phi_2 \colon E_1 \to E_2$ be normalized isogenies. Then $\phi_1 = \phi_2$. In other words, for any fixed E_1 and E_2, there exists at most one normalized isogeny $E_1 \to E_2$.*

Proof. We follow the proof given in [1]. Let $\hat{\phi}_1$ be the dual of ϕ_1. Then we have two endomorphisms $\psi_1 = \hat{\phi}_1 \circ \phi_1$ and $\psi_2 = \hat{\phi}_1 \circ \phi_2$ of E_1. Notice also that $c_{\psi_1} = c_{\hat{\phi}_1} = c_{\psi_2}$, as ϕ_1 and ϕ_2 are normalized. By Lemma 1, we see $c_{\psi_1 - \psi_2} = 0$. This means that the endomorphism $\psi_1 - \psi_2$ is constant, and therefore 0. So $\psi_1 = \psi_2$, and likewise $\phi_1 = \phi_2$. ☐

Proposition 5. *Let E/k, E_1/k, E_2/k be elliptic curves. Suppose $\phi_1 \colon E \to E_1$ and $\phi_2 \colon E \to E_2$ are normalized isogenies with the same kernel. Then $E_1 = E_2$ and $\phi_1 = \phi_2$.*

Proof. By Proposition 12.12 (and Remark 12.13) of [9], we know there is an isomorphism $\psi : E_1 \to E_2$ defined over k such that $\phi_2 = \psi \circ \phi_1$. By Lemma 2, $c_\psi = 1$. As $\mathrm{char}(k) \nmid 6$, we may assume our curves have Weierstrass equations of the form $y^2 = x^3 + ax + b$. Thus, this isomorphism ψ must be of the form $\psi(x, y) = (u^2 x, u^3 y)$, for some $u \in k^*$. It is easy to see that $c_\psi = u$, hence we must have $u = 1$, and ψ is the identity map. The result is now immediate. ☐

Remark: For arbitrary characteristic, we must allow general Weierstrass equations of the form $y^2 + a_1xy + a_3y = x^3 + a_2x^2 + a_4x + a_6$. In this case, a version of Proposition 5 is still true, but we must add the condition that both of the isogenies ϕ_1 and ϕ_2 fix a_1, a_2, and a_3 (meaning these coefficients are the same for all three curves). The isogeny given by Vélu's formula has this property, so given any normalized isogeny ϕ, with kernel $\mathcal{F}$, if ϕ fixes a_1, a_2, and a_3, then ϕ is equal to the isogeny given by Vélu's formula.

2.3. **An exact sequence with Vélu's isogeny.** In this section, we consider elliptic curves over finite fields F_q, $q = p^f$, of characteristic $p > 3$. Let $E_1/\mathbb{F}_q$ be an elliptic curve, and suppose $P \in E_1(\mathbb{F}_q)$ is a point of exact order m, where $(q, m) = 1$. Let τ_P denote the m-isogeny $\tau_P \colon E_1 \to E_2$ whose kernel is exactly $\langle P \rangle$, as given by Vélu's formula. The subgroup $\tau_P(E_1(\mathbb{F}_q)) \leq E_2(\mathbb{F}_q)$ determines an isogeny $\psi \colon E_2 \to E_3$, also given by Vélu's formula.

Theorem 6. *The codomain of ψ is exactly the curve E_1. Moreover, $\psi(E_2(\mathbb{F}_q)) = \langle P \rangle$. Equivalently, we have the following exact sequence of abelian groups:*

$$\begin{tikzcd}\{\infty\} \arrow[r] & \langle P \rangle \arrow[r] & E_1(\mathbb{F}_q) \arrow[r, "\tau_P"] & E_2(\mathbb{F}_q) \arrow[r, "\psi"] & \langle P \rangle \arrow[r] & \{\infty\}\end{tikzcd} .$$

Proof. We compare two isogenies, α and β. Define $\alpha = \psi \circ \tau_P$, and $\beta = 1 - \pi_q$, where π_q denotes the Frobenius endomorphism $(x, y) \mapsto (x^q, y^q)$. Both maps are separable. This is clear for α, as it is a composition of separable isogenies; β is known to be separable by [9, Prop. 2.29]. Moreover, they are both normalized. For α, this follows from Corollary 3 and the fact that Vélu's formula provides a normalized isogeny. For β, this follows from the fact ([6, III.5.5]) that $(s + t\pi_q)^*\omega = s\omega$ for any integers $s, t \in \mathbb{Z}$. These isogenies must therefore be equal by Proposition 5. Thus, the codomain of α (and hence ψ) is the codomain of β, which is E_1.

Let $\mathcal{F} = \psi(E_2(\mathbb{F}_q))$. It remains only to show that $\mathcal{F} = \langle P \rangle$. Certainly, we have the exact sequence

$$\begin{tikzcd}\{\infty\} \arrow[r] & \langle P \rangle \arrow[r] & E_1(\mathbb{F}_q) \arrow[r, "\tau_P"] & E_2(\mathbb{F}_q) \arrow[r, "\psi"] & \mathcal{F} \arrow[r] & \{\infty\}\end{tikzcd} .$$

As E_1 and E_2 are isogenous, the middle two groups have the same cardinality, and so by exactness we must have $\#\mathcal{F} = m$. Let ϕ be the isogeny on E_2 with kernel $\mathcal{F}$ which is given by Vélu's formula. Essentially repeating the argument of the previous paragraph (this time considering $1 - \pi_q$ on E_2), we see that $\phi \circ \psi$ has codomain E_2. Thus, both $\tau_P \circ \psi$ and $\phi \circ \psi$ are normalized isogenies $E_2 \to E_2$ with kernel

$E_2(\mathbb{F}_q)$. By Proposition 4, they must be the same. Hence $\tau_P = \phi$, and so $\mathcal{F} = \langle P \rangle$. $\square$

3. A character sum determined by an isogeny

We now extend some of the results of [4] (where P is always a point of order $m = 2$) to general m. Let $E_1/\mathbb{F}_q$ be an elliptic curve, and $P \in E_1(\mathbb{F}_q)$ a point of exact order m, with $(q, m) = 1$. Let $\tau_P \colon E_1 \to E_2$ be the isogeny with kernel $\langle P \rangle$ given by Vélu's formula. By Theorem 6, the quotient $E_2(\mathbb{F}_q)/\tau_P(E_1(\mathbb{F}_q))$ is isomorphic to the cyclic group $\langle P \rangle$. Choose $Q \in E_2(\mathbb{F}_q)$ such that $[Q]$ generates the quotient. When m is prime, we may take any $Q \in E_2(\mathbb{F}_q) - \tau_P(E_1(\mathbb{F}_q))$. Let $\zeta = \zeta_m$ be a primitive m-th root of unity. We define a character on $E_2(\mathbb{F}_q)$, determined by τ_P and $[Q]$:

$$\chi_P(R) = \zeta^j, \text{ for } R - jQ \in \tau_P(E_1(\mathbb{F}_q)).$$

We define an associated character sum on the points of E_2:

$$S_P := \sum_{\substack{R \in E_2(\mathbb{F}_q) \\ R \neq \infty}} \chi_P(R) x_R. \tag{6}$$

Note that S_P depends on the choice of $[Q]$. However, as $\psi(E_2(\mathbb{F}_q)) = \langle P \rangle$, we may always make a "preferred" choice of $[Q]$ such that $\psi(Q) = P$. Under this convention, we will see that S_P depends only on P. In fact, S_P depends only on the cyclic subgroup $\langle P \rangle$, and not the particular generator.

Theorem 7. *Let $\psi \colon E_2 \to E_1$ be the normalized isogeny with kernel $\tau_P(E_1(\mathbb{F}_q))$ given by Vélu. Then*

$$S_P = \sum_{j=1}^{m-1} \zeta^j x_{\psi(jQ)}.$$

Moreover, if Q is chosen so that $\psi(Q) = P$, then

$$S_P = \sum_{j=1}^{m-1} \zeta^j x_{jP}.$$

Note the right hand side is a character sum of the x-coordinates of points on E_1, while the left hand side is defined over points on E_2.

Proof. Let $\hat{\tau}_P$ be the dual isogeny to τ_P, so that $\hat{\tau}_P \circ \tau_P = [m]$ on E_1 and $\tau_P \circ \hat{\tau}_P = [m]$ on E_2, where $[m]$ is the multiplication by m map. Notice that if $U = mR$ for some $R \in E_2(\mathbb{F}_q)$, then $U = \tau_P(\hat{\tau}_P(R))$, and hence $U \in \tau_P(E_1(\mathbb{F}_p))$.

We first examine when $m = 2k$ is even. Let $\mathcal{T} = \tau_P(E_1(\mathbb{F}_q))$. Then we have

$$S_P = \sum_{\substack{R\in\mathcal{T}\\ R\neq\infty}} x_R + \sum_{j=1}^{k-1}\left(\sum_{R\in\mathcal{T}}(\zeta^j+\zeta^{-j})x_{jQ+R}\right) + \sum_{R\in\mathcal{T}}\zeta^k x_{kQ+R}.$$

Using Vélu's formula (5) (and noting $\zeta^k = -1$), we have

$$\begin{aligned}
S_P &= -x_{kQ} + \sum_{\substack{R\in\mathcal{T}\\ R\neq\infty}}(x_R - x_{kQ+R}) + \sum_{j=1}^{k-1}\left(\sum_{R\in\mathcal{T}}(\zeta^j+\zeta^{-j})x_{jQ+R}\right)\\
&= -x_{\psi(kQ)} + \sum_{j=1}^{k-1}\left(\sum_{R\in\mathcal{T}}(\zeta^j+\zeta^{-j})x_{jQ+R}\right)\\
&= -x_{\psi(kQ)} + \sum_{j=1}^{k-1}(\zeta^j+\zeta^{-j})\sum_{R\in\mathcal{T}} x_{jQ+R}\\
&= -x_{\psi(kQ)} + \sum_{j=1}^{k-1}(\zeta^j+\zeta^{-j})\left(\sum_{R\in\mathcal{T}} x_{jQ+R} - \sum_{\substack{R\in\mathcal{T}\\ R\neq\infty}} x_R + \sum_{\substack{R\in\mathcal{T}\\ R\neq\infty}} x_R\right)\\
&= -x_{\psi(kQ)} + \sum_{j=1}^{k-1}(\zeta^j+\zeta^{-j})\left(x_{\psi(jQ)} + \sum_{\substack{R\in\mathcal{T}\\ R\neq\infty}} x_R\right)\\
&= -x_{\psi(kQ)} + \sum_{j=1}^{k-1}(\zeta^j+\zeta^{-j})x_{\psi(jQ)} + \sum_{j=1}^{k-1}(\zeta^j+\zeta^{-j})\sum_{\substack{R\in\mathcal{T}\\ R\neq\infty}} x_R\\
&= -x_{\psi(kQ)} - (\zeta^k+1)\sum_{\substack{R\in\mathcal{T}\\ R\neq\infty}} x_R + \sum_{j=1}^{k-1}(\zeta^j+\zeta^{-j})x_{\psi(jQ)}\\
&= -x_{\psi(kQ)} + \sum_{j=1}^{k-1}(\zeta^j+\zeta^{-j})x_{\psi(jQ)} = \sum_{j=1}^{m-1}\zeta^j x_{\psi(jQ)}.
\end{aligned}$$

The case when m is odd is proven similarly. This proves the first claim. The second claim follows immediately. □

4. Low degree isogenies

We now apply the results of Theorem 7 to specific low degree isogenies. Throughout this section we set $q = p$, so that $\mathbb{F}_q = \mathbb{F}_p$, with p

prime. In the sections that follow, we use parameterizations of elliptic curves with a point of exact order $m \leq 8$. Such parameterizations are well-known, and can be found in [2] or [3] for instance. For our purposes, we will use the parametrization given in [5] for the case $m = 4$, and the parametrizations found by Kubert in [3] for $m = 3$, $5 \leq m \leq 8$. In principle these results could be extended to any value of m for which such parameterizations exist, e.g., when $X_1(m)$ has genus 0.

In each case, $E_1/\mathbb{F}_p$ is an elliptic curve, $P \in E_1(\mathbb{F}_p)$ has exact order m, $\tau_P \colon E_1 \to E_2$ is the isogeny given by Vélu whose kernel is $\langle P \rangle$, and $Q \in E_2(\mathbb{F}_p)$ is chosen so that $[Q]$ generates $E_2(\mathbb{F}_p)/\tau_P(E_1(\mathbb{F}_p))$. We always assume $(p, m) = 1$, and let $\zeta \in \overline{\mathbb{F}}_p$ denote a primitive m-th root of unity.

4.1. **The case $m = 2$.**

When $m = 2$, Theorem 7 becomes

$$S_P = -x_{\psi(Q)}.$$

If P is a point of order 2, then by a suitable change of variables we may assume that $P = (0, 0)$. As $Q \notin \tau_P(E_1(\mathbb{F}_p)) = \ker \psi$, by Theorem 6, we must have $\psi(Q) = P$. So $S_P = 0$. This is an alternate way to show that $S_{\tau,p}$, as defined in (4), satisfies $S_{\tau,p} \equiv 0 \pmod{p}$ when $a = 0$.

4.2. **The case $m = 3$.**

If $P = (a, b)$, then we claim $S_P = -a$. This follows from Theorem 7, as Q must satisfy $3Q = \infty$. Thus, $\psi(2Q) = -\psi(Q)$, and

$$\begin{aligned} S_P &= \zeta x_{\psi(Q)} + \zeta^2 x_{\psi(2Q)} \\ &= \zeta x_{\psi(Q)} + \zeta^2 x_{-\psi(Q)} = -x_{\psi(Q)} = -a. \end{aligned}$$

Notice that in this case S_P is independent of the choice of Q.

It is well known that any elliptic curve over $\mathbb{F}_p$ with an $\mathbb{F}_p$-rational point P of order 3 is isomorphic to a curve of the form

$$E_1 : y^2 + \alpha xy + \beta y = x^3, \qquad P = (0, 0)$$

where $\alpha, \beta \in \mathbb{F}_p$, $\alpha^3(\beta^3 - 27\alpha) \neq 0$. Since $\psi(Q) \in \{P, 2P\}$ and $2P = (0, -\beta)$, we have $S_P = 0$.

4.3. **The case $m = 4$.**

Let $P = (a, b)$, and write $2P = (c, d)$. By Theorem 7 (as $3Q = -Q$),

$$S_P = \zeta x_{\psi(Q)} + \zeta^2 x_{\psi(2Q)} + \zeta^3 x_{\psi(3Q)} = -x_{2\psi(Q)} = -c.$$

We see again that S_P is independent of the choice of Q.

Every elliptic curve over $\mathbb{F}_p$ with an $\mathbb{F}_p$-rational point of order 4 is isomorphic to a curve of the form

$$E_1 : y^2 = x^3 + (1 - 2\alpha)x^2 + \alpha^2 x, \qquad P = (\alpha, \alpha)$$

with $\alpha \in \mathbb{F}_p$, $\alpha \neq 0, \frac{1}{4}$ in $\mathbb{F}_p$. On this curve, $2P = (0,0)$, and so $S_P = 0$.

4.4. **The case** $m = 5$.

Write $P = (a,b)$, $2P = (c,d)$. By Theorem 7:

$$\begin{aligned} S_P &= \zeta x_{\psi(Q)} + \zeta^2 x_{\psi(2Q)} + \zeta^3 x_{\psi(3Q)} + \zeta^4 x_{\psi(4Q)} \\ &= \lambda x_{\psi(Q)} + \bar{\lambda} x_{2\psi(Q)} = \lambda a + \bar{\lambda} c, \end{aligned}$$

where $\lambda = \zeta + \zeta^{-1}$, $\bar{\lambda} = \zeta^2 + \zeta^{-2}$ are the roots of $x^2 + x - 1$ in $\mathbb{F}_p(\sqrt{5})$. The value of S_P lies in $\mathbb{F}_p(\sqrt{5})$, and is a root of

$$f(x) = x^2 + (a+c)x - a^2 + 3ac - c^2 \in \mathbb{F}_p[x].$$

This value depends on Q, although $f(x)$ does not.

Every elliptic curve over $\mathbb{F}_p$ with an $\mathbb{F}_p$-rational point of order 5 is isomorphic to a curve of the form

$$E_1 : y^2 + (1+\alpha)xy + \alpha y = x^3 + \alpha x^2, \qquad P = (0,0),$$

with $\alpha \in \mathbb{F}_p$ and $\alpha(\alpha^2 + 11\alpha - 1) \neq 0$. Regardless of the choice of Q, the set $\{a, c\} = \{0, -\alpha\}$, and so the possible values of S_P are $-\lambda\alpha$, $-\bar{\lambda}\alpha$, the roots of $x^2 - \alpha x - \alpha^2$. Although S_P does not vanish, note that there is always a choice of Q such that $S_P = s + t\sqrt{5}$, with $s + t = 0$.

4.5. **The case** $m = 6$.

We have

$$\begin{aligned} S_P &= \zeta x_{\psi(Q)} + \zeta^2 x_{\psi(2Q)} + \zeta^3 x_{\psi(3Q)} + \zeta^4 x_{\psi(4Q)} + \zeta^5 x_{\psi(5Q)} \\ &= (\zeta + \zeta^{-1}) x_{\psi(Q)} + (\zeta^2 + \zeta^{-2}) x_{2\psi(Q)} - x_{3\psi(Q)} \\ &= x_{\psi(Q)} - x_{2\psi(Q)} - x_{3\psi(Q)}. \end{aligned}$$

Note that $\psi(Q) = \pm P$, and S_P gives the same result for either value. So S_P is independent of Q.

Every elliptic curve over $\mathbb{F}_p$ with an $\mathbb{F}_p$-rational point of order 6 is isomorphic to a curve of the form

$$E_1 : y^2 + (1-\alpha)xy - (\alpha + \alpha^2)y = x^3 - (\alpha + \alpha^2)x^2, \qquad P = (0,0),$$

with $\alpha \in \mathbb{F}_p$, $\alpha \neq 0, -1, -\frac{1}{9}$. We find $S_P = -\alpha(\alpha + 2)$.

4.6. **The case** $m = 7$.

Writing $\lambda_k = \zeta^k + \zeta^{-k}$, we have

$$S_P = \lambda_1 x_{\psi(Q)} + \lambda_2 x_{\psi(2Q)} + \lambda_3 x_{\psi(3Q)},$$

which is dependent on the choice of Q. However, analogous to the $m = 5$ case, there is a cubic polynomial $f \in \mathbb{F}_p[x]$ whose roots give all possible values for S_P, and whose coefficients are polynomials in the x-coordinates of the points in $\langle P \rangle$. If we let a, b, c denote the x-coordinates of P, $2P$, $3P$, respectively, then

$$\begin{aligned} f(x) = x^3 &+ (a+b+c)x^2 + (3ab + 3ac + 3bc - 2a^2 - 2b^2 - 2c^2)x \\ &+ (4ab^2 + 4bc^2 + 4ca^2 - 3a^2b - 3b^2c - 3c^2a - a^3 - b^3 - c^3). \end{aligned}$$

Every elliptic curve over $\mathbb{F}_p$ with an $\mathbb{F}_p$-rational point of order 7 is isomorphic to a curve of the form

$$E_1\colon y^2+(1+\alpha-\alpha^2)xy-\alpha^2(\alpha-1)y=x^3-\alpha^2(\alpha-1), \qquad P=(0,0),$$

where $\alpha\in\mathbb{F}_p$, $\alpha(\alpha-1)(\alpha^3-8\alpha^2+5\alpha+1)\neq 0$. In this case, we simplify $f(x)$ above and find $S_P=\alpha(\alpha-1)T$, where T is a root of

$$g(x)=x^3+(\alpha+1)x^2-(2\alpha^2-3\alpha+2)x-(\alpha^3+3\alpha^2-4\alpha+1),$$

with the particular root depending on the choice of Q.

4.7. **The case $m=8$.** We find

$$S_P=-x_{\psi(4Q)}+\sqrt{2}(x_{\psi(Q)}-x_{\psi(3Q)}),$$

which is dependent on the choice of Q. Let a, c, d, denote the x-coordinates of P, $3P$, $4P$, respectively. Regardless of the choice of Q, S_P is a root of

$$f(x)=x^2+2dx+(d^2-2(a-c)^2)\in\mathbb{F}_p[x].$$

Every elliptic curve over $\mathbb{F}_p$ with an $\mathbb{F}_p$-rational point of order 8 is isomorphic to a curve of the form

$$E_1: y^2-(2\alpha^2-4\alpha+1)xy-\alpha\beta y=x^3-\beta x^2, \qquad P=(0,0),$$

where $\alpha\in\mathbb{F}_p$, $\alpha(\alpha-1)(\alpha-\frac{1}{2})(\alpha^2-\alpha+\frac{1}{8})\neq 0$, and $\beta=(2\alpha-1)(\alpha-1)\alpha^2$. We find

$$S_P=\alpha(1-\alpha)\big(\alpha^2\pm(2\alpha-1)\sqrt{2}\big),$$

dependent on the choice of Q.

Acknowledgments

We acknowledge the contribution of SAGE [7], which facilitated the construction of examples which were helpful in discovering the main theorems of this work. The first author would also like to thank Daniel Shumow for conversations which led to a better understanding of normalized isogenies. This research was partially supported by the National Institute of Standards and Technology.

References

[1] J. González. On the division polynomials of elliptic curves. *Rev. R. Acad. Cienc. Exactas Fís. Nat. (Esp.)*, 94(3):377–381, 2000. Contributions to the algorithmic study of problems of arithmetic moduli (Spanish).

[2] T. Hadano. Elliptic curves with a torsion point. *Nagoya Math. J.*, 66:99–108, 1977.

[3] D. S. Kubert. Universal bounds on the torsion of elliptic curves. *Proc. London Math. Soc. (3)*, 33(2):193–237, 1976.

[4] C. McLeman and C. Rasmussen. Class number formulas via 2-isogenies of elliptic curves. *Bulletin of the London Mathematical Society*, 2012.

[5] R. Rezaeian Farashahi, D. Moody, and H. Wu. Isomorphism classes of Edwards curves over finite fields. *Finite Fields Appl.*, 18(3):597–612, 2012.

[6] J. H. Silverman. *The arithmetic of elliptic curves*, volume 106 of *Graduate Texts in Mathematics*. Springer, Dordrecht, second edition, 2009.

[7] W. Stein et al. *Sage Mathematics Software (Version 4.7)*. The Sage Development Team, 2011. `http://www.sagemath.org`.

[8] J. Vélu. Isogénies entre courbes elliptiques. *C. R. Acad. Sci. Paris Sér. A-B*, 273:A238–A241, 1971.

[9] L. C. Washington. *Elliptic curves*. Discrete Mathematics and its Applications (Boca Raton). Chapman & Hall/CRC, Boca Raton, FL, second edition, 2008. Number theory and cryptography.